ELSEVIER International Journal of Mass Spectrometry and Ion Processes 136 (1994) 215

AUTHOR INDEX

Abe, T., 85

Baptista, G.B., 107
Barros Leite, C.V., 107
Begemann, F., 91
Birkinshaw, K., 71
Bowie, J.H., 143
Bozek, J.D., 55
Bradford, A.M., 143
Burgers, P.C., 191

Campbell, E.E.B., 209
Cisper, M.E., 119

da Silveira, E.F., 107
de Petris, G., 155

Eiden, G.C., 119
Ernstberger, B., 101

Franklin, A.M., 25

Garrett, A.W., 119
Grebner, T.L., 101
Gross, M.L., 143

Hemberger, P.H., 119
Hiroki, S., 85

Jeronymo, J.M.F., 107

Langstaff, D.P., 71
Liu, W.G., 181

Mann, M., 167
McGibbon, G.A., 191
Murakami, Y., 85

Neusser, H.J., 101
Nguyen, M.T., 45
Nogar, N.S., 119

Occhiucci, G., 155
Operti, L., 25
Ott, U., 91

Pepi, F., 155
Prinzbach, H., 209

Ray, K.B., 107
Richter, S., 91

Saito, N., 55
Schweikert, E.A., 107
Splendore, M., 25
Suzuki, I.H., 55

Takayama, M., 35
Tellgmann, R., 209
Terlouw, J.K., 191
Todd, J.F.J., 25

Vaglio, G.A., 25
Vollmer, D.L., 143

Wahl, F., 209
Waugh, R.J., 143
Weitzel, K.-M., 1
Wilm, M.S., 167

Xiao, Y.K., 181

Zhou, Y.M., 181

ELSEVIER International Journal of Mass Spectrometry and Ion Processes 136 (1994) 217–219

SUBJECT INDEX

Ab initio calculations
Hydrogen cyanide loss from $[CH_5N_2]^+$ cations: 1,2-elimination versus Beckmann rearrangement, 45
Tunneling RRKM calculations for the H_2 loss reaction from ethane ions on an ab initio potential energy surface, 1
Atomic weight
Precise measurement of the isotopic composition of cerium and its atomic weight, 181

Beckmann rearrangement
Hydrogen cyanide loss from $[CH_5N_2]^+$ cations: 1,2-elimination versus Beckmann rearrangement, 45
Branching ratio
Variation in ion-pair yields from CF_2Cl_2 photoexcited in the inner-shell excitation regions, 55

Carbenes
The imidic acids H–N=C(H)–OH and CH_3–N=C(H)–OH and their tautomeric carbenes H_2N–$\ddot{C}$–OH and CH_3–N(H)–$\ddot{C}$–OH: stable species in the gas phase formed by one-electron reduction of their cations, 191
Cation affinity
Formation and collision-induced dissociation of adduct ions $[matrix + C]^+$ (C = Li, Na, Cs and NH_4) produced under fast atom bombardment conditions, 35
Cerium
Precise measurement of the isotopic composition of cerium and its atomic weight, 181
CF_2Cl_2
Variation in ion-pair yields from CF_2Cl_2 photoexcited in the inner-shell excitation regions, 55
Channeltrons
Multiple ion counting in isotope abundance mass spectrometry, 91
Coincidence counting
Coincidence counting analysis of secondary ions emitted from a phenylalanine target under multi-charged MeV carbon bombardment, 107
Collision-induced dissociation
Collision-induced dissociations of deprotonated dipeptide methyl esters containing H, alkyl or benzyl α side-chains, 143
Formation and collision-induced dissociation of adduct ions $[matrix + C]^+$ (C = Li, Na, Cs and NH_4) produced under fast atom bombardment conditions, 35
The imidic acids H–C=C(H)–OH and CH_3–N=C(H)–OH and their tautomeric carbenes H_2N–$\ddot{C}$–OH and CH_3–N(H)–$\ddot{C}$–OH: stable species in the gas phase formed by one-electron reduction of their cations, 191
Coordination structure
Formation and collision-induced dissociation of adduct ions $[matrix + C]^+$ (C = Li, Na, Cs and NH_4) produced under fast atom bombardment conditions, 35
Correlated ion emission
Coincidence counting analysis of secondary ions emitted from a phenylalanine target under multi-charged MeV carbon bombardment, 107

Detector array
Resolving power of a discrete electrode detector array, 71
Diazapropylium cations
Hydrogen cyanide loss from $[CH_5N_2]^+$ cations: 1,2-elimination versus Beckmann rearrangement, 45
Dichlorodibenzo-*p*-dioxin
The ionization energy of 2,8-dichlorodibenzo-*p*-dioxin by resonance-enhanced two-photon ionization, 101
Dipeptide methyl esters
Collision-induced dissociations of deprotonated dipeptide methyl esters containing H, alkyl or benzyl α side-chains, 143

Dodecahedranes
Fullerene formation via laser-induced oligomerisation of 1,6-dibromo-dodecahedrane ($C_{20}H_{18}Br_2$), 209
Dynamic trapping
An improved method for capturing laser desorbed ions in an ion trap mass spectrometer: dynamic r.f. trapping, 119

Electrospray
Electrospray and Taylor-Cone theory, Dole's beam of macromolecules at last? 167
1,2-Elimination
Hydrogen cyanide loss from $[CH_5N_2]^+$ cations: 1,2-elimination versus Beckmann rearrangement, 45

Fringing field
Influence of the fringing field length on the separated $^4He/D_2$ peak shape of a high-resolution quadrupole mass spectrometer, 85
Fullerenes
Fullerene formation via laser-induced oligomerisation of 1,6-dibromo-dodecahedrane ($C_{20}H_{18}Br_2$), 209

Germane
Investigation of the complex reactions of SiH_4 and GeH_4 in the ion trap using dynamically programmed scanning, 25

Heats of formation
Hydrogen cyanide loss from $[CH_5N_2]^+$ cations: 1,2-elimination versus Beckmann rearrangement, 45
High resolution
Influence of the fringing field length on the separated $^4He/D_2$ peak shape of a high-resolution quadrupole mass spectrometer, 85
Hydrazonium cation
Hydrogen cyanide loss from $[CH_5N_2]^+$ cations: 1,2-elimination versus Beckmann rearrangement, 45

Imidic acids
The imidic acids H–N=C(H)–OH and CH_3–N=C(H)–OH and their tautomeric carbenes H_2N–$\ddot{C}$–OH and CH_3–N(H)–$\ddot{C}$–OH: stable species in the gas phase formed by one-electron reduction of their cations, 191
Inner-shell excitation
Variation in ion-pair yields from CF_2Cl_2 photoexcited in the inner-shell excitation regions, 55
Integrated circuit
Resolving power of a discrete electrode detector array, 71

Ion desorption
Coincidence counting analysis of secondary ions emitted from a phenylalanine target under multi-charged MeV carbon bombardment, 107
Ion injection
An improved method for capturing laser desorbed ions in an ion trap mass spectrometer: dynamic r.f. trapping, 119
Ion/molecule reactions
Investigation of the complex reactions of SiH_4 and GeH_4 in the ion trap using dynamically programmed scanning, 25
Ion trap
An improved method for capturing laser desorbed ions in an ion trap mass spectrometer: dynamic r.f. trapping, 119
Ion trap mass spectrometry
Investigation of the complex reactions of SiH_4 and GeH_4 in the ion trap using dynamically programmed scanning, 25
Isotope abundance mass spectrometry
Multiple ion counting in isotope abundance mass spectrometry, 91
Isotope ratio measurement
Precise measurement of the isotopic composition of cerium and its atomic weight, 181

Kinetics
Investigation of the complex reactions of SiH_4 and GeH_4 in the ion trap using dynamically programmed scanning, 25

Laser desorption
An improved method for capturing laser desorbed ions in an ion trap mass spectrometer: dynamic r.f. trapping, 119
Laser-induced oligomerisation
Fullerene formation via laser-induced oligomerisation of 1,6-dibromo-dodecahedrane ($C_{20}H_{18}Br_2$), 209

Mass spectrometry
An improved method for capturing laser desorbed ions in an ion trap mass spectrometer: dynamic r.f. trapping, 119
MS–MS
Electrospray and Taylor-Cone theory, Dole's beam of macromolecules at last? 167
Multiple ion counting
Multiple ion counting in isotope abundance mass spectrometry, 91

One-electron reduction
The imidic acids H–N=C(H)–OH and CH_3–N=C(H)–OH and their tautomeric carbenes H_2N–$\ddot{C}$–OH and CH_3–N(H)–$\ddot{C}$–OH: stable species in the gas phase formed by one-electron reduction of their cations, 191
Organic/metallic adduct
Formation and collision-induced dissociation of adduct ions $[matrix + C]^+$ (C = Li, Na, Cs and NH_4) produced under fast atom bombardment conditions, 35

Partial yield spectra
Variation in ion-pair yields from CF_2Cl_2 photoexcited in the inner-shell excitation regions, 55
Peptides
Electrospray and Taylor-Cone theory, Dole's beam of macromolecules at last? 167
Phosphorous and phosphonic acid
A mass spectrometric study of gaseous $H_4PO_3^+$ and $H_2PO_3^-$ ions, 155
Phosphorus polyanions
A mass spectrometric study of gaseous $H_4PO_3^+$ and $H_2PO_3^-$ ions, 155
Proteins
Electrospray and Taylor-Cone theory, Dole's beam of macromolecules at last? 167
Proton affinity
A mass spectrometric study of gaseous $H_4PO_3^+$ and $H_2PO_3^-$ ions, 155
Formation and collision-induced dissociation of adduct ions $[matrix + C]^+$ (C = Li, Na, Cs and NH_4) produced under fast atom bombardment conditions, 35

Quadrupole mass spectrometer
Influence of the fringing field length on the separated $^4He/D_2$ peak shape of a high-resolution quadrupole mass spectrometer, 85

Rate constants
Tunneling RRKM calculations for the H_2 loss reaction from ethane ions on an ab initio potential energy surface, 1
Reaction mechanism
Tunneling RRKM calculations for the H_2 loss reaction from ethane ions on an ab initio potential energy surface, 1
Resolving power
Resolving power of a discrete electrode detector array, 71
Resonance-enhanced multiphoton ionization
The ionization energy of 2,8-dichlorodibenzo-*p*-dioxin by resonance-enhanced two-photon ionization, 101

Selected photoion–photoion coincidence
Variation in ion-pair yields from CF_2Cl_2 photoexcited in the inner-shell excitation regions, 55
Silane
Investigation of the complex reactions of SiH_4 and GeH_4 in the ion trap using dynamically programmed scanning, 25
Silicon
Resolving power of a discrete electrode detector array, 71

Tautomerism
A mass spectrometric study of gaseous $H_4PO_3^+$ and $H_2PO_3^-$ ions, 155
Taylor Cone
Electrospray and Taylor-Cone theory, Dole's beam of macromolecules at last? 167
Time-of-flight mass spectrometry
Fullerene formation via laser-induced oligomerisation of 1,6-dibromo-dodecahedrane ($C_{20}H_{18}Br_2$), 209
The ionization energy of 2,8-dichlorodibenzo-*p*-dioxin by resonance-enhanced two-photon ionization, 101
Transition state
Tunneling RRKM calculations for the H_2 loss reaction from ethane ions or an ab initio potential energy surface, 1
Tunneling
Tunneling RRKM calculations for the H_2 loss reaction from ethane ions on an ab initio potential energy surface, 1